MINNESOTA

CANCER BURDEN



In 2004, the American Cancer Society estimates that

- 1,368,030 new cancer cases will be diagnosed in the United States, including 22,720 in Minnesota.
- 563,700 cancer deaths will occur in the United States, including 9,360 in Minnesota.

The average annual age-adjusted death rate for cancer per 100,000 persons²

Minnesota: 187.5 National: 199.8

LUNG CANCER

In 2004, the American Cancer Society estimates that1

- 2,580 new cases of lung cancer will be diagnosed among men and women in Minnesota.
- 2,380 men and women will die of lung cancer in Minnesota.

The CDC National Center for Health Statistics provides the following death rates:

The average annual age-adjusted death rates for lung cancer per 100,000 persons, by race, 1997–2001^{2,3}

| | Minnesota | National |
|------------------------------|-----------|----------|
| Overall | 47.1 | 56.2 |
| White | 46.6 | 56.2 |
| Black | 77.7 | 65.2 |
| Hispanic | 36.3 | 24.9 |
| Asian/Pacific Islander | 29.2 | 28.2 |
| American Indian/Alaska Nativ | e 91.7 | 36.3 |

BREAST CANCER

In 2004, the American Cancer Society estimates that¹

- 3,610 new cases of breast cancer will be diagnosed among women in Minnesota.
- 670 women will die of breast cancer in Minnesota.

The CDC National Center for Health Statistics provides the following death rates:

The average annual age-adjusted death rates for breast cancer per 100,000 persons, by race, 1997–2001^{2,3}

| | Minnesota | National |
|------------------------------|-----------|----------|
| Overall | 25.9 | 27.0 |
| White | 26.0 | 26.4 |
| Black | 33.5 | 35.4 |
| Hispanic | 21.7 | 17.2 |
| Asian/Pacific Islander | - | 12.6 |
| American Indian/Alaska Nativ | e - | 13.6 |

Visit http://www.cdc.gov/cancer/npcr/index.htm — *United States Cancer Statistics* — for more information on top cancer sites by geographic area, race, and gender.

COLORECTAL CANCER

In 2004, the American Cancer Society estimates that¹

- 2,200 new cases of colorectal cancer will be diagnosed among men and women in Minnesota.
- 850 men and women will die of colorectal cancer in Minnesota.

The CDC National Center for Health Statistics provides the following death rates:

The average annual age-adjusted death rates for colorectal cancer per 100,000 persons, by race, 1997–2001^{2,3}

| | Minnesota | National |
|------------------------------|-----------|----------|
| Overall | 19.0 | 20.8 |
| White | 19.0 | 20.3 |
| Black | 22.9 | 28.3 |
| Hispanic | - | 14.2 |
| Asian/Pacific Islander | 11.7 | 13.0 |
| American Indian/Alaska Nativ | e 28.2 | 13.9 |

PROSTATE CANCER

In 2004, the American Cancer Society estimates that¹

- 4,230 new cases of prostate cancer will be diagnosed among men in Minnesota.
- 550 men will die of prostate cancer in Minnesota.

The CDC National Center for Health Statistics provides the following death rates:

The average annual age-adjusted death rates for prostate cancer per 100,000 men, by race, 1997–2001^{2,3}

| | Minnesota | National |
|------------------------------|-----------|----------|
| Overall | 32.9 | 31.5 |
| White | 32.6 | 28.8 |
| Black | 68.6 | 70.4 |
| Hispanic | - | 23.6 |
| Asian/Pacific Islander | - | 13.0 |
| American Indian/Alaska Nativ | e - | 20.2 |

¹ Source: American Cancer Society Facts and Figures, 2004. Estimates exclude more than a million cases of basal and squamous cell skin cancers and *in situ* cancers, except urinary bladder, that will be diagnosed in 2004. Lung cancer rates include bronchus cancer. State death totals were rounded to nearest 10.

² Source: Centers for Disease Control and Prevention (CDC) National Center for Health Statistics, vital statistics data, underlying cause of death, 1997–2001. Death rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population.

³ Hyphens represent suppression of rates when there were 75,000 or fewer persons in the denominator or 20 or fewer deaths in the numerator.